

DOI: 10.21767/2471-9943.100041

Vitamin B₁₂ Status after Right Hemicolectomy in Bowel Cancer Patients: A Feasibility Study

Williams A^{1#}, Carter J^{2#}, Basu P¹, Hughes L¹, Nair D¹, Williams EM³ and Edwards P^{1*}

¹Department of Surgery, Nevill Hall Hospital, Aneurin Bevan University Health Board, Abergavenny, UK

²Department of Clinical Biochemistry, Nevill Hall Hospital, Aneurin Bevan University Health Board, Abergavenny, UK

³Faculty of Life Sciences and Education, University of South Wales, Newport, UK

Williams A and Carter J contributed equally to this work.

*Corresponding author: Paul Edwards, Department of Surgery, Nevill Hall Hospital, Aneurin Bevan University Health Board, Abergavenny, UK, Tel: 01873732412; E-mail: Paul.Edwards3@wales.nhs.uk

Received: June 22, 2017; Accepted: July 05, 2017; Published: July 11, 2017

Citation: Williams A, Carter J, Basu P, Hughes L, Nair D, et al. (2017) Vitamin B₁₂ Status after Right Hemicolectomy in Bowel Cancer Patients: A Feasibility Study. Colorec Cancer 3:2.

Copyright: © 2017 Williams A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Introduction: Terminal ileum resection in patients with Crohn's disease results in vitamin B₁₂ deficiency and these patients are routinely given B₁₂ supplementation to prevent anemia and neuropsychiatric disease. The effect of right hemicolectomy on B₁₂ status in cancer patients has not previously been studied. Total serum B₁₂ is an unreliable test. Second-line diagnostic tests measure methylmalonic acid (MMA) and homocysteine (tHcy) are also used. An assay measuring an active form of B₁₂ is available for routine use. This observational study investigated whether B₁₂ deficiency exists in patients after a right hemicolectomy for cancer.

Methods: Patients (n=28; age range=52-88 years) who had undergone a right hemicolectomy for cancer and been in remission for two years were identified from a database and recruited to an outpatient clinic along with an age-matched control group (n=27; age range=52-89 years). Vitamin B₁₂ status in both patients and controls was assessed by the measurement of vitamin B₁₂ (total and active), MMA and tHcy.

Results: Concentration of tHcy was significantly higher ($p<0.05$) in the surgical group (14.0 μ mol/L) compared with the controls (11.1 μ mol/L). There were no significant differences in total and active B₁₂, folate and MMA concentrations between the two groups ($p>0.05$).

Discussion: Right hemicolectomy for malignancy is associated with an increase in tHcy concentrations although a larger study is required to determine the B₁₂ status in post-operative right hemicolectomy patients.

Keywords: Right hemicolectomy; Vitamin B₁₂ deficiency; MMA; tHcy; Cancer; Crohn's disease

Introduction

Cancer Research UK reports bowel cancer as the fourth most common cancer for both men and women in the UK, with 41,112 bowel cancer cases in the UK, and 2,272 in Wales in 2013 [1]. Sixty- three per cent of bowel cancer patients had surgical intervention in 2015 [2] with some undergoing a right hemicolectomy with various degrees of the terminal ileum being resected.

There has been no study to investigate vitamin B₁₂ deficiency in people undergoing right hemicolectomy for malignancy however those who have had the same procedure for Crohn's disease are prescribed B₁₂ supplementation because of recognised deficiencies. Crohn's ileal resections of less than 20 cm in length have variable consequences on B₁₂ status [3] although patients who have had more than 20 cm of terminal ileum excised almost invariably have vitamin B₁₂ deficiency [4].

Vitamin B₁₂ plays a key role in DNA and cell metabolism and long term deficiency can cause severe and irreversible damage. Effects of B₁₂ deficiency include fatigue, bone marrow suppression, neurological features including stroke, and risk of cardiomyopathy [5-8]. It has been recognised that B₁₂ deficiency can cause symptoms of psychosis, mania, memory impairment, fatigue, irritation and personality changes. Deficiency can be treated with vitamin B₁₂ supplementation, which if given for life has the potential to improve well-being and longevity.

Vitamin B₁₂ (cobalamin) is bound to two proteins; transcobalamin (TC) and haptocorrin (HC). The transcobalamin-vitamin B₁₂ complex, also known as holotranscobalamin (HolotC), contains the biologically available B₁₂. HolotC mediates the uptake of B₁₂ by all cells utilising specific receptors whereas 80% of B₁₂ which is bound to HC is metabolically inert [8,9]. Measurement of total B₁₂ (cobalamin) in serum by immunoassay is the most widely available test for assessing B₁₂ status. However, this test

measures both the active and inactive forms and has a low sensitivity. Currently methylmalonic acid (MMA) and homocysteine (tHcy) are recommended as second line tests if total B₁₂ results are in the indeterminate range, as they are considered to be more sensitive tests although tHcy is less specific. Both MMA and tHcy depend on B₁₂ as a cofactor for their metabolism. An increase in MMA or total tHcy in folate replete individuals indicates B₁₂ deficiency in tissue [7,10]. The metabolically active B₁₂ can now be measured by an immunoassay and recent evidence suggests that this is a more reliable marker of impaired vitamin B₁₂ status than total B₁₂ [7,10]. However, it is also recommended that MMA is used as a confirmatory test if the result is in the indeterminate range.

In this present study, we measure both total and active B₁₂, tHcy and MMA in addition to other variables to determine B₁₂ status in patients who have undergone right hemicolectomies for cancer and in age-matched controls (controls).

Table 1 Descriptive characteristics for all participants (n=55) and for controls (n=27) and patient (n=28) subgroups.

	Total cohort	Controls	Patients	P
n (%)	55 (100%)	27 (49%)	28 (51%)	
Age (years)	75 (52-89)	77 (52-89)	74 (52-88)	0.933
Gender (M:F)	40:15	22:5	18:10	-
Total B ₁₂ (ng/L)	317 (125-895) ^d	324 (195-895) ^b	295 (125-652) ^b	0.243
Active B ₁₂ (pmol/L)	65.3 (23.4-257.0) ^a	69.3 (23.4-257.0)	63.7 (27.5-170.1) ^a	0.782
Folate (μg/L)	5.8 (1.5-20.0) ^d	5.8 (1.5-20.0) ^b	5.7 (2.3-16.0) ^b	0.902
MMA (μmol/L)	0.29 (0.16-2.80)	0.28 (0.21-0.81)	0.33 (0.16-2.80)	0.619
Homocysteine (μmol/L)	12.6 (6.5-38.0)	11.1 (6.9-34.5)	14 (6.5-38.0)	0.037
Haemoglobin (g/L)	136 ± 19 ^a	133 ± (24) ^a	139 ± (15)	0.256
MCV (fL)	89.0 (79-112) ^a	89 (80-112) ^a	90 (79-104)	0.549
Creatinine (μmol/L)	82 (59-148) ^c	77 (61-119) ^b	84 (59-148) ^a	0.223
eGFR (mL/min/1.73 m ²) (mean ± SD)	73 ± 17 ^c	78 ± 15 ^b	68 ± 17 ^a	0.040
TI Excised (cms)	-	-	80 (60-400)	-
Days post resection			1747 (770-2812) ^a	

Note: continuous variables are expressed as median (range) unless stated otherwise. tHcy, homocysteine; MMA, methylmalonic acid; eGFR, estimated glomerular filtration rate; MCV, mean cell volume. Missing number of samples are shown, a: n-1, b: n- 2, c: n-3, and d: n-4.

Laboratory analyses

Blood samples were collected, transported to the laboratory within 30 minutes and analysed on the same day with the exception of plasma tHcy, plasma MMA, and active vitamin B₁₂, where samples were stored at -80°C for up to one month prior to analysis. For plasma tHcy, samples were separated and frozen within 30 minutes of venepuncture.

Serum total B₁₂, folate, creatinine and tHcy, were analysed on the Abbott ARCHITECT ci8200 analyser (Abbott Diagnostics, Berkshire, UK). Full blood count parameters including haemoglobin and mean cell volume (MCV) were performed by a SYSMEX XE-2100 analyser (SYSMEX, Kobe, Japan). Plasma

Materials and Methods

Recruitment of participants

Adults treated with a right hemicolectomy for gastrointestinal cancer at Nevill Hall Hospital who had been in remission for two years or more were identified from an internal database and invited to join the study. Potential participants were excluded with the following conditions: already receiving vitamin B₁₂ supplementation, pregnant, anemic or unable to give written consent. The study gained ethical approval from the NHS South Central Hampshire B Research ethics committee (REC No 15/SC/0756 PR). All participants gave written informed consent. Twenty-eight participants attended a clinic where blood samples were taken along with twenty-seven age-matched controls (Table 1).

concentrations of MMA were determined by GC-MS. Serum active B₁₂ was analysed by an Abbott chemiluminescent microparticle immunoassay (CMIA) on the Abbott ARCHITECT ci8200 analyser. Within-batch coefficients of variation (CVs) were 4.1% and 5.3% at concentrations of 15.8 pmol/L and 46.8 pmol/L, respectively. Between-batch CVs were 7.6% and 5.7% at concentrations of 15.8 pmol/L and 46.8 pmol/L, respectively. Glomerular filtration rate was estimated using a four-variable MDRD equation. For the purpose of this study, renal disease was defined as estimated GFR (eGFR) <60 mL/min/1.73 m² [11]. Local reference ranges were used for all analytes with the exception of the reference range for active B₁₂ [12].

Statistical Analysis

Data were analysed using statistical software (Analyse-It, Leeds UK and SigmaPlot V13 Systat Ltd, UK). Most of the data were not normally distributed and were expressed as median (range) unless stated otherwise. Non-parametric Mann-Whitney U-test was used for comparison between patients and controls. For normally distributed data the unpaired T-test was used. Spearman rank analysis was used to test for univariate relationships between markers of vitamin B₁₂ status with clinical variables. A *p* value of <0.05 was considered statistically significant. Agreement between different markers of B₁₂ status was undertaken using the weighted kappa test for agreement: kappa statistic (κ) <0.2 was considered a poor strength of agreement.

Results

Eleven of 26 (42%) patients and 10 of 26 (38%) controls had total B₁₂ concentrations within the indeterminate range (180-300 ng/L) (**Table 1**). In comparison, a slightly higher proportion of patients (56% (15 of 27)) and controls (48% (13 of 27)) had active B₁₂ concentrations in the indeterminate range (25-70 pmol/L). However, 3 of 26 (12%) patients demonstrated total B₁₂ deficiency (<180 ng/L) compared with none in the control group. Conversely, active B₁₂ results revealed that 1 of 27 controls (4%) was deficient (<25 pmol/L) compared with none in the patient group. Taken together, there was a high proportion of patients who were not replete in B₁₂; 56% and 53% for active B₁₂ total B₁₂, respectively. A slightly lower proportion of controls were not replete in B₁₂; 50% for active B₁₂ and 40% for total B₁₂.

Of those study participants (patients and controls) who were not replete in active B₁₂ (<70 pmol/L), only 9 of 30 (30%) and 8 of 30 (27%) had raised MMA (>0.42 μ mol/L) and tHcy (>16.0 μ mol/L) concentrations, respectively. Study participants who were not replete in total B₁₂ (<300 ng/L), 4 of 24 (17%) and 5 of 24 (21%) had raised MMA (>0.42 μ mol/L) and tHcy (>16.0 μ mol/L) concentrations, respectively.

(>16.0 μ mol/L) concentrations, respectively. Thus, agreement between total B₁₂ and MMA (κ =0.02), active B₁₂ and MMA (κ =0.07), tHcy and total B₁₂ and tHcy (κ =0.01), active B₁₂ and tHcy (κ = 0.02) for the diagnosis of B₁₂ deficiency was poor in this population. Of those with deficient total B₁₂ results (<180 ng/L; n=3 patients), active B₁₂ results were all in the indeterminate range or above, one MMA and two tHcy results were elevated, and one tHcy result was at the upper end of the reference range. In contrast, only one control had a deficient active B₁₂ result (<25 pmol/L). However total B₁₂ was in the indeterminate range and MMA and tHcy concentrations were both normal.

Plasma homocysteine concentration (14.0 μ mol/L) was significantly higher in patients compared with controls (11.1 μ mol/L) (*p*=0.037). Ten of 28 (36%) patients exhibited an elevated tHcy concentration (>16.0 μ mol/L) compared with 4 of 27 (15%) of controls. eGFR was significantly lower in patients (68 mL/min/1.73 m^2) compared with controls (78 mL/min/1.73 m^2) (*p*=0.04). However, only seven of the patients and four of the controls had eGFR <60 mL/min/1.73 m^2 . There were no significant differences (*p*>0.05) in total or active B₁₂, MMA, haemoglobin (Hb) and mean cell volume (MCV) between the two groups. The prevalence for abnormal MMA (>0.42 μ mol/L) in patients and controls was 21% (6 of 28) and 19% (5 of 27), respectively.

Univariate analyses for the patient subgroup (**Table 2**) revealed that tHcy correlated negatively with folate and eGFR and positively with MMA. However, in the controls, tHcy only correlated negatively with folate (**Table 3**). In patients, total B₁₂ only correlated positively with active B₁₂ compared with total B₁₂ in control participants which correlated positively with active B₁₂, folate and MCV. In patients, active B₁₂ correlated positively with total B₁₂ and negatively with MMA, and Hb. In the controls active B₁₂ correlated negatively with folate and positively with total B₁₂. Due to the small study numbers, the independent effect of variables on B₁₂ status was not analyzed.

Table 2 Correlation between vitamin B₁₂ status markers and clinical variables in patients (n=28).

	tHcy		Total B ₁₂		Active B ₁₂		MMA	
	<i>r</i> _s	<i>p</i>	<i>r</i> _s	<i>P</i>	<i>r</i> _s	<i>P</i>	<i>r</i> _s	<i>P</i>
tHcy	-	-	-0.27	0.19	-0.25	0.20	0.43	0.022
Total B ₁₂	-0.27	0.19	-	-	0.49	0.01	-0.16	0.42
Active B ₁₂	-0.25	0.20	0.49	0.01	-	-	-0.55	0.003
MMA	0.43	0.02	-0.16	0.42	-0.55	0.003	-	-
Folate	-0.74	<0.0001	0.02	0.90	0.24	0.24	-0.45	0.02
Hb	-0.06	0.76	-0.32	0.11	-0.44	0.02	0.32	0.10
eGFR	-0.52	0.005	-0.25	0.22	-0.07	0.72	-0.12	0.53
MCV	0.19	0.34	0.19	0.34	0.14	0.48	0.04	0.82
TI	0.17	0.38	-0.20	0.33	-0.04	0.83	0.02	0.93
Resection	0.06	0.75	0.10	0.64	0.10	0.61	-0.12	0.54

Note: rs, correlation coefficient.; tHcy, homocysteine; MMA, methylmalonic acid; Hb, haemoglobin; eGFR, estimated glomerular filtration rate; MCV, mean cell volume; TI, terminal ileum length; Resection, days post resection.

Table 3 Correlation between vitamin B₁₂ status markers and clinical variables in control group (n=27).

	tHcy		Total B ₁₂		Active B ₁₂		MMA	
	r _s	P	r _s	p	r _s	p	r _s	P
tHcy	-		-0.26	0.20	-0.32	0.10	0.27	0.18
Total B ₁₂	-0.26	0.20	-	-	0.51	0.01	0.24	0.25
Active B ₁₂	-0.32	0.10	0.51	0.01	-	-	-0.13	0.52
MMA	0.27	0.18	0.24	0.25	-0.13	0.52	-	-
Folate	-0.62	0.001	0.43	0.03	0.44	0.03	0.12	0.57
Hb	0.19	0.34	0.05	0.98	-0.00	0.99	-0.03	0.89
eGFR	-0.36	0.08	0.35	0.10	0.34	0.10	-0.08	0.69
MCV	-0.20	0.32	0.46	0.02	0.36	0.07	0.07	0.74

Note: rs, correlation coefficient; tHcy, homocysteine; MMA, methylmalonic acid; Hb, haemoglobin; eGFR, estimated glomerular filtration rate; MCV, mean cell volume.

Discussion

This is the first study to assess B₁₂ status in patients who have undergone a right hemicolectomy for cancer. The study has demonstrated that tHcy concentrations were mildly elevated in patients compared with controls. Over twice as many patients had abnormal tHcy concentrations (>16.0 μmol/L) compared with controls. While vitamin B₁₂ deficiency results in an elevated plasma tHcy, and is a sensitive marker of vitamin B₁₂ deficiency, it is not specific to vitamin B₁₂ deficiency and is a nonspecific marker for many diseases including cardiovascular diseases and dementia [13-15]. Recent studies suggest that mildly elevated concentrations of tHcy may be an indicator of atherosclerosis and cardiovascular disease [14]. However, mildly elevated concentrations (<50 μmol/L) frequently occur in the general population and are influenced by many factors, including folate deficiency, vitamin B₆ deficiency poor renal function, hypothyroidism, lifestyle factors (smoking, alcoholism, diet) age and gender, drug therapy and certain genetic polymorphisms. The main predictors of elevated plasma tHcy concentrations are low folate, vitamin B₁₂ status or renal impairment [13,14]. Although an inverse relationship exists between tHcy and folate in both patients and controls in this study, all participants, apart from one patient and one control, were defined as folate replete (>2.4 μg/L). Similarly, this study demonstrated an inverse relationship between tHcy and eGFR. Hyperhomocysteinaemia (<50 μmol/L) is known to occur when eGFR falls below 60 mL/min/1.73 m² [16]. Furthermore, it has been demonstrated that renal impairment causes an increase in tHcy and MMA concentrations in an aged population (≥ 65 years) [17]. In the present study, six of the ten patients with raised tHcy concentrations exhibited concurrent impaired renal function (eGFR <60 mL/min/1.73 m²) whereas four patients had normal renal function (eGFR >60 mL/min/1.73 m²). However, the assessment of renal function (eGFR <60 mL/min/ 1.73 m²) in this study is based on a single measurement

and it is recommended that diagnosis of chronic kidney disease is recorded over at least three months [18]. On repeat testing some participants with eGFR <60 mL/min/1.73 m² may exhibit a higher eGFR and vice versa. Notably, the median age of our population is 77 years and thus an effect on renal function is likely which may in turn compromise the interpretation of tHcy and MMA results. The independent effect of renal function on the measurement of tHcy and MMA was not investigated.

The thyroid function and vitamin B₆ status of this group were not investigated and cannot be ruled out as causes. It is possible that these patients also have cardiovascular disease but details of this co-morbidity were not recorded at recruitment and was not part of the exclusion criteria. Further work is necessary to establish the cause of mildly elevated tHcy in this patient group.

Although the study did not show significant differences in active B₁₂ concentrations between patients and controls there was an inverse relationship between active B₁₂ and MMA within the patient cohort which was not observed amongst controls. Due to the small numbers of participants in this study, the independent effect of variables on MMA and B₁₂ concentrations was not explored.

This study highlights the difficulty in defining B₁₂ status using various circulating markers. The total B₁₂ and active B₁₂ results both demonstrate that a large proportion of patients and controls fall within the indeterminate range. According to recent British Journal of Haematology guidelines, patients with a reduced total B₁₂ and normal tHcy and MMA do not require any further investigations and are not classified as B₁₂ deficient [19]. In the current study, further testing using the recommended second-line investigations, MMA or tHcy, revealed that only a quarter of the study participants with indeterminate active B₁₂ results and only a fifth with indeterminate total B₁₂ results, may be B₁₂ deficient. Although many studies recommend these second line investigations

there is currently not a gold standard test available for B_{12} status. Recently, active B_{12} has been shown to be the best marker of B_{12} status in older people (n=700; mean age=81 years) when compared with red cell cobalamin [7]. Furthermore, in this study the indeterminate range was defined as 20-30 pmol/L and anything above that was defined as normal. If these cut-offs are applied to these data then all study participants would be defined as replete despite having elevated MMA and tHcy concentrations, thereby highlighting the importance of establishing local reference ranges due to variations between populations and methodologies. For the present study, the reference ranges for active B_{12} recommended by the referral laboratory in London have been used, which may not represent the population in South Wales. It would be useful to establish a local reference range for active B_{12} . Thus, further work is necessary before the B_{12} status can be accurately determined in our local population and also in cancer patients who have undergone right hemicolectomy.

Of the participants classified as B_{12} deficient (determined by either total or active B_{12}) none were found to have macrocytosis and/or anemia. Similarly, a Finnish study investigating the prevalence of B_{12} deficiency in an elderly population (n=1048; age range: 65-100 years) demonstrated that anemia or macrocytosis did not predict B_{12} deficiency and only two of their 97 subjects with B_{12} deficiency would have been diagnosed based on these observations [20]. This raises the question of the sensitivity of this approach in the early diagnosis of B_{12} deficiency.

The small sample size in this study does not allow a detailed analysis of B_{12} status as originally anticipated. These results demonstrate that a high proportion of patients (54%) and controls (48%) are not replete in vitamin B_{12} . Whether this is a true representation of the local population needs further investigation and a further larger study would allow a better stratification of study participants into those who are vitamin B_{12} deficient or replete. The population (600,000) served by Aneurin Bevan University Health Board come from both rural and suburban areas, with a mix of socioeconomic status. Those living in rural and lower socioeconomic areas may have a limited access to healthcare and as a consequence are more likely to be vitamin B_{12} deficient.

In conclusion, this study demonstrates that there are subtle differences between the controls and patients who have undergone a right hemicolectomy for cancer. We recommend that vitamin B_{12} status is investigated further using the aforementioned recommended laboratory markers for tissue vitamin B_{12} stores in order to understand how terminal ileum excision alters this essential regulatory system. Replenishing vitamin B_{12} may improve short and long-term outcomes of patients that have had a right hemicolectomy.

Acknowledgements

The authors are grateful to the laboratory staff in the Blood Sciences Department at Nevill Hall Hospital, Abergavenny for processing the samples for all analytes apart from MMA and

tHcy. They are also grateful to the laboratory staff in the Department of Biochemistry at University Hospital of Wales, Cardiff for processing the samples for MMA and tHcy analyses and to Abbott Diagnostics for providing the active B_{12} assay. The study received financial support from the R&D Department at ABUHB.

References

1. Cancer Research UK [<http://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/bowel-cancer/incidence.>] Accessed on February 2017.
2. National Bowel Cancer Audit Report (2016) The Association of Colorectal Cancer of Great Britain and Ireland.
3. Duerksen DR, Fallows G, Bernstein CN (2006) Vitamin B_{12} malabsorption in patients with limited ileal resection. Nutrition 22: 1210-1213.
4. Thompson WG, Wrathell E (1977) The relation between ileal resection and Vitamin B_{12} absorption. Can J Surg 20: 461-464.
5. Behrend C, Jeppesen PB, Mortensen PB (1995) Vitamin B_{12} absorption after ileorectal anastomosis for Crohn's Disease. Eur J Gastroenterol Hepatol 7: 397-400.
6. Domstad PA, Choy YC, Kim EE, DeLand FH (1981) Reliability of the Dual Isotope Schilling test for the diagnosis of Pernicious Anemia or Malabsorption Syndromes. Am J Clin Pathol 75: 723-726.
7. Valente E, Scott JM, Per-Magne U, Cunningham C, Casey M, et al. (2011) Diagnostic accuracy of Holotranscobalamin, methylmalonic acid, serum cobalamin and other indicators of tissue Vitamin B_{12} status in the elderly. Clin Chem 57: 856-863.
8. Spence JD (2016) Metabolic vitamin B_{12} deficiency: a missed opportunity to prevent dementia and stroke. Nutr Res 36: 109-116.
9. Hunt A, Harrington D, Robinson S (2014) Vitamin B_{12} deficiency. BMJ 349: 5226.
10. Nexo E, Horrmann-Lucke E (2011) Holotranscobalamin, a marker of vitamin B_{12} status: analytical aspects and clinical utility. Am J Clin Nutr 94: 359S-365S.
11. Levey AS, Greene T, Kusek J (2000) A simplified equation to predict glomerular filtration rate from serum creatinine. J Am Soc Nephrol 11: 155A.
12. Viapath (2017) [www.viapath.co.uk] Accessed on February 2017.
13. Refsum H, Smith AD, Ueland PM, Nexo E, Clarke R, et al. (2004) Facts and recommendations about total homocysteine determinations: an expert opinion. Clin Chem 50: 3-32.
14. Moat SJ (2008) Plasma total homocysteine: instigator or indicator of cardiovascular disease? Ann Clin Biochem 45: 345-348.
15. McCaddon A, Husdon P, Davies G, Hughes A, William JH, et al. (2001) Homocysteine and cognitive decline in healthy elderly. Dement Geriatr Cogn Disord 12: 309-313.
16. Van GC (2006) Why is homocysteine elevated in renal failure and what can be expected from homocysteine-lowering? Nephrol Dial Transplant 21: 1161-1166.
17. Loikas S, Koskinen P, Irljala K, Loppönen M, Isoaho R, et al. (2007) Renal impairment compromises the use of total homocysteine

and methylmalonic acid but not total vitamin B12 and holotranscobalamin in screening for vitamin B12 deficiency in the aged. *Clin Chem Lab Med* 45: 197-220.

18. Levey AS, Eckardt KU, Tsukamoto Y, Levin A, Coresh J, et al. (2005) Definition and classification of chronic kidney disease: a position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int* 67: 2089-2100.

19. Devalia V, Hamilton MS, Molloy AM (2014) Guidelines for the diagnosis and treatment of cobalamin and folate disorders. *Br J Haem* 166: 496-513.

20. Loikas S, Koskinen P, Irljala K, Löppönen M, Isoaho R, et al. (2007) Vitamin B12 deficiency in the aged: a population-based study. *Age Ageing* 36: 177-183.